arXiv:1409.7503v3 [math.FA] 4 Feb 2015

APPROXIMATE BIPROJECTIVITY AND ϕ-BIFLATNESS OF CERTAIN BANACH ALGEBRAS

A. SAHAMI AND A. POURABBAS

Abstract. In the first part of the paper, we investigate the approximate biprojectivity of some Banach algebras related to the locally compact groups. We show that a Segal algebra $S(G)$ is approximately biprojective if and only if G is compact. Also for every continuous weight w, we show that $L^1(G,w)$ is approximately biprojective if and only if G is compact.

In the second part, we study ϕ-biflatness of some Banach algebras, where ϕ is a Banach algebra character. We show that if $S(G)$ is ϕ-biflat, then G is an amenable group for every character ϕ. Finally we show that ϕ-biflatness of $L^1(G)^{**}$ implies the amenability of G.

1. Introduction and Preliminaries

The concepts of ϕ-biflatness, ϕ-biprojectivity, ϕ-Johnson amenability and other related concepts were introduced and studied in [17]. The studies include determining when the various classes of Banach algebras are, or are not ϕ-biflat or ϕ-biprojective. It was shown in [17] that $L^1(G)$ is ϕ-biflat if and only if G is an amenable group and the Fourier algebra $\mathcal{A}(G)$ is ϕ-biprojective if and only if G is a discrete group.

Recently the concepts of approximate biprojectivity and approximate biflatness have been studied by Zhang [20] and Samei *et al.* [19], respectively. Samei *et al.* in [19] studied approximate biflatness of Segal algebras and Fourier algebras and they showed that the Segal algebra $S(G)$ is pseudo-contractible if and only if G is compact. Note that the pseudo-contractility of Banach algebras implies the approximate biprojectivity [5, Proposition 3.8], that is, the approximate biprojectivity is a weaker notion than the pseudo-contractility, for more details see [5].

Motivated by these results, in this paper we extend [19, Theorem 3.5] or [2, Theorem 5.3] and we show that Segal algebra $S(G)$ is approximately biprojective if and only if G is compact. The group algebra $L^1(G)$ is biprojective if and only if G is compact, see [6, Theorem 5.13]. Here we extend this result, we show that the weighted group algebra $L^1(G,w)$ is approximately biprojective if and only if G is compact for every continuous weight w on G. We show that if Segal algebra $S(G)$ is ϕ-biflat, then G is amenable, where ϕ is any character on $S(G)$ and if $L^1(G)^{**}$ is $\tilde{\phi}$-biflat, then G is amenable, where $\tilde{\phi}$ is an extension of character ϕ on $L^1(G)$.

We remark some standard notations and definitions that we shall need in this paper. Let A be a Banach algebra. If X is a Banach A-bimodule, then X^* is also a Banach A-bimodule via the following actions

$$(a\cdot f)(x)=f(x\cdot a),\quad (f\cdot a)(x)=f(a\cdot x)\qquad (a\in A, x\in X, f\in X^*).$$

2010 *Mathematics Subject Classification.* Primary 43A07, 43A20, Secondary 46H05.

Key words and phrases. Beurling algebras, Segal algebras, Approximate biprojectivity, ϕ-biflat.

Throughout, the character space of A is denoted by $\Delta(A)$, that is, all non-zero multiplicative linear functionals on A. Let $\phi \in \Delta(A)$. Then ϕ has a unique extension $\tilde{\phi} \in \Delta(A^{**})$ which is defined by $\tilde{\phi}(F) = F(\phi)$ for every $F \in A^{**}$.

Let A be a Banach algebra. The projective tensor product $A \otimes_p A$ is a Banach A-bimodule via the following actions

$$a \cdot (b \otimes c) = ab \otimes c, \ (b \otimes c) \cdot a = b \otimes ca \qquad (a, b, c \in A).$$

The product morphism $\pi_A : A \otimes_p A \to A$ is specified by $\pi_A(a \otimes b) = ab$ for every $a, b \in A$.

Let G be a locally compact group. The Fourier algebra on G is denoted by $\mathcal{A}(G)$. It is well-known that the character space $\Delta(\mathcal{A}(G))$ consists of all point evaluation maps $\phi_t : \mathcal{A}(G) \to \mathbb{C}$ such that $\phi_t(f) = f(t)$ for each $f \in \mathcal{A}(G)$, see [3].

We also remind some concepts of Banach homology which we shall need in this paper. A Banach algebra A is called biprojective, if there exists a bounded A-bimodule morphism $\rho : A \to A \otimes_p A$ such that ρ is a right inverse for π_A [6]. We recall that A is an approximately biprojective Banach algebra if there exists a net of bounded A-bimodule morphism $(\rho_\alpha) : A \to A \otimes_p A$ such that $\pi_A \circ \rho_\alpha(a) \to a$ for each $a \in A$, see [20]. A Banach algebra A is called ϕ-biflat for every $\phi \in \Delta(A)$, if there exists a bounded A-bimodule morphism $\rho : A \to (A \otimes_p A)^{**}$ such that $\tilde{\phi} \circ \pi_A^{**} \circ \rho(a) = \phi(a)$ for every $a \in A$, [17]. Also A is called left ϕ-amenable (left ϕ-contractible) if there exists an element $m \in A^{**}$ $(m \in A)$ such that $am = \phi(a)m$ and $\tilde{\phi}(m) = 1$ $(\phi(m) = 1)$ for every $a \in A$, respectively. For more details on left ϕ-amenability and left ϕ-contractibility see [9] and [12], respectively.

Following theorem is given by authors in [18]. They characterized approximate biprojectivity of some semigroup algebras. We apply this theorem in order to characterize approximate biprojectivity of algebras related to the locally compact groups.

Theorem 1.1. [18, Theorem 3.9] *Let A be an approximately biprojective Banach algebra with a left approximate identity (right approximate identity) and let $\phi \in \Delta(A)$. Then A is left ϕ-contractible(right ϕ-contractible), respectively.*

2. Approximate biprojectivity

In this section we improve [19, Theorem 3.5] or [2, Theorem 5.3] and [6, Theorem 5.13] concerning approximate biprojectivity of some Banach algebras related to the locally compact groups.

We remind that a Banach algebra A is called pseudo-contractible if there is a not necessarily bounded net $(m_\alpha)_\alpha$ in $A \otimes_p A$ such that $a \cdot m_\alpha = m_\alpha \cdot a$ and $\pi_A(m_\alpha)a \to a$ for each $a \in A$. For the fundamental details of the pseudo-contractibility readers are referred to [5] and [2].

Now we consider Segal algebras on a locally compact group. As we see in [14] a Segal algebra $S(G)$ on a locally compact group G is a dense left ideal of $L^1(G)$ that satisfies the following conditions:

(i) $S(G)$ is a Banach space with respect to a norm $||\cdot||_S$ satisfying $||\cdot||_{L^1} \leq ||\cdot||_S$.

(ii) For $f \in S(G)$ and $y \in G$, $\delta_y * f \in S(G)$ and the map $y \mapsto \delta_y * f$ is continuous. Also $||\delta_y * f||_S = ||f||_S$, for $f \in S(G)$ and $y \in G$.

With the norm $||\cdot||_S$ and the convolution product, $S(G)$ is a Banach algebra and we have the following inequality

$$||f * g||_S \leq ||f||_{L^1}||g||_S \qquad f \in L^1(G),\ g \in S(G).$$

$S(G)$ on a locally compact group G always has a left approximate identity and it is never amenable unless it is $L^1(G)$ itself and G is amenable. A Segal algebra is called symmetric if for $f \in S(G)$ and $y \in G$, $f * \delta_y \in S(G)$ and the map $y \mapsto f * \delta_y$ is continuous and also $||f * \delta_y||_S = ||f||_S$. By [14, Proposition 1, page 19] a symmetric Segal algebra is an ideal in $L^1(G)$ which

$$||g * f||_S \leq ||f||_{L^1}||g||_S \qquad f \in L^1(G),\, g \in S(G).$$

Note that $\Delta(S(G)) = \{\phi_{|_{S(G)}} | \phi \in \Delta(L^1(G))\}$ and ϕ_0 (the augmentation character on $L^1(G)$) induces a character on $S(G)$ still denoted by ϕ_0 [1, Lemma 2.2].

Samei *et al.* in [19, Theorem 3.5] and Choi *et al.* in [2, Theorem 5.3] showed that $S(G)$ is pseudo-contractible if and only if G is compact. As approximate biprojectivity is weaker notion than pseudo-contractibility in the following theorem we extend this result.

Theorem 2.1. *Let G be a locally compact group. Then $S(G)$ is approximately biprojective if and only if G is compact.*

Proof. Let $S(G)$ be approximately biprojective. Since $S(G)$ has a left approximate identity, Theorem 1.1 shows that $S(G)$ is left ϕ_0-contractible, hence by [12, Theorem 2.1] there exists an element $m \in S(G)$ such that $a * m = \phi_0(a)m$ and $\phi_0(m) = 1$ for every $a \in S(G)$. Since $S(G)$ is dense in $L^1(G)$, it is easy to see that $a * m = \phi_0(a)m$ and $\phi_0(m) = 1$ for every $a \in L^1(G)$. Now apply [12, Theorem 6.1] to show that G is compact.

Converse is clear by [19, Theorem 3.5] or [2, Theorem 5.3]. □

Theorem 2.2. *Let G be an SIN group. If $S(G)\otimes_p S(G)$ is approximately biprojective, then G is compact.*

Proof. The main result of [11] asserts that, if G is an SIN group, then $S(G)$ has a central approximate identity, say $(e_\alpha)_{\alpha\in I}$. Since $S(G) \otimes_p S(G)$ is approximately biprojective, there exists a net

$$(\rho_\beta)_{\beta\in\Theta} : S(G) \otimes_p S(G) \to (S(G) \otimes_p S(G)) \otimes_p (S(G) \otimes_p S(G))$$

of continuous $S(G) \otimes_p S(G)$-bimodule morphism such that $\pi_{S(G)\otimes_p S(G)} \circ \rho_\beta(x) \to x$, for every $x \in S(G) \otimes_p S(G)$. Consider $n_\alpha = e_\alpha \otimes e_\alpha$, it is easy to see that for every $x \in S(G) \otimes_p S(G)$ we have $xn_\alpha = n_\alpha x$ and $\phi\otimes\phi(n_\alpha) = \phi\otimes\phi(e_\alpha\otimes e_\alpha) = \phi(e_\alpha)\phi(e_\alpha) \to 1$, where $\phi \in \Delta(S(G))$. Define $m_\alpha^\beta = \rho_\beta(n_\alpha)$. Then it is easy to see that $x \cdot m_\alpha^\beta = m_\alpha^\beta \cdot x$ for every $x \in S(G) \otimes_p S(G)$. Also

$$\begin{aligned} \lim_\alpha \lim_\beta \phi \otimes \phi \circ \pi_{S(G)\otimes_p S(G)}(m_\alpha^\beta) - 1 &= \lim_\alpha \lim_\beta \phi \otimes \phi \circ \pi_{S(G)\otimes_p S(G)} \circ \rho_\beta(n_\alpha) - 1 \\ &= \lim_\alpha \phi \otimes \phi(n_\alpha) - 1 \\ &= \lim_\alpha \phi(e_\alpha)^2 - 1 = 0. \end{aligned} \tag{2.1}$$

Set $E = I \times \Theta^I$, where Θ^I is the set of all functions from I into Θ. Consider the product ordering on E as follows

$$(\alpha, \beta) \leq_E (\alpha', \beta') \Leftrightarrow \alpha \leq_I \alpha', \beta \leq_{\Theta^I} \beta' \qquad (\alpha, \alpha' \in I, \quad \beta, \beta' \in \Theta^I),$$

here $\beta \leq_{\Theta^I} \beta'$ means that $\beta(d) \leq_\Theta \beta'(d)$ for each $d \in I$. Suppose that $\gamma = (\alpha, \beta_\alpha) \in E$ and $m_\gamma = \rho_{\beta_\alpha}(n_\alpha) \in (S(G) \otimes_p S(G)) \otimes_p (S(G) \otimes_p S(G))$. Now using the iterated limit theorem [10, page 69] in (2.1) we obtain

$$\phi \otimes \phi \circ \pi_{S(G)\otimes_p S(G)}(m_\gamma) \to 1$$

and similarly we obtain $x \cdot m_\gamma = m_\gamma \cdot x$ for every $x \in S(G) \otimes_p S(G)$. By using the same argument as in the proof of [17, Proposition 2.2] one can show that $S(G) \otimes_p S(G)$ is left $\phi \otimes \phi$-contractible. Hence [12, Theorem 3.14] shows that $S(G)$ is left ϕ-contractible. So $L^1(G)$ is left ϕ-contractible. Applying [12, Theorem 6.1] G must be compact. □

Let G be a locally compact group. A real-valued function w on G is said to be a weight function if it has the following properties:

(i) $w(x) \geq 1 \quad (x \in G)$,
(ii) $w(xy) \geq w(x)w(y) \quad (x, y \in G)$,
(iii) w is measurable and locally bounded.

We form the Banach space

$$L^1(G, w) = \left\{ f : G \to \mathbb{C} : \ fw \in L^1(G) \right\}.$$

Then $L^1(G, w)$, with the convolution product, is a Banach algebra and is called Beurling algebra. See [15] for further information on Beurling algebras.

Helemskii [6, Theorem 5.13] showed that the group algebra $L^1(G)$ is biprojective if and only if G is compact. At the following theorem we extend this result.

Theorem 2.3. *Let G be a locally compact group and let w be a continuous weight on G. Then $L^1(G, w)$ is approximately biprojective if and only if G is compact.*

Proof. Let $L^1(G, w)$ be approximately biprojective. Since $L^1(G, w)$ has a left approximate approximate identity [15, Propostion 3.7.7], by Theorem 1.1 $L^1(G, w)$ is left ϕ-contractible for every $\phi \in \Delta(L^1(G, w))$ even for the augmentation character ϕ_0 which is specified by

$$\phi_0(f) = \int_G f(x)dx.$$

By [12, Theorem 2.1] there exists an element $m \in L^1(G, w)$ such that $a * m = \phi_0(a)m$ and $\phi_0(m) = 1$ for every $a \in L^1(G, w)$. Pick $f \in L^1(G, w)$ such that $\phi_0(f) = 1$. We have

$$\delta_g * m = \phi_0(f)\delta_g * m = \delta_g * (f * m) = (\delta_g * f) * m = \phi_0(\delta_g * f)m = \phi_0(f)m = m,$$

which shows that m is a constant function in $L^1(G, w)$, so we can assume that $1 \in L^1(G, w)$. Since $w(g) \geq 1$ for every $g \in G$, we have

$$|G| = \int_G 1dg \leq \int_G w(g)dg < \infty.$$

Now apply [7, Theorem 15.9] to show that G is compact.

For converse, using the same argument as in [6, Theorem 5.13], it is easy to see that $L^1(G, w)$ is biprojective, so $L^1(G, w)$ is approximately biprojective. □

Proposition 2.4. *Let G be a locally compact group and let A be a unital Banach algebra with $\Delta(A) \neq \emptyset$. If $A \otimes_p L^1(G)$ is approximately biprojective, then G is compact and A is approximately biprojective. Converse holds if A is biprojective.*

Proof. Suppose that $B = A \otimes_p L^1(G)$ is approximate biprojective. It is easy to see that $(e_A \otimes e_\alpha)$ is an approximate identity for B, where e_A is an identity for A and (e_α) is a bounded approximate identity for $L^1(G)$. Let $\psi \in \Delta(A)$ and $\phi \in \Delta(L^1(G))$. Then Theorem 1.1 implies that B is left $\psi \otimes \phi$-contractible.

By [12, Theorem 3.14] $L^1(G)$ is left ϕ-contractible which implies that G is compact, see [12, Theorem 6.1].

Let $\rho : G \to \mathbb{C}$ be a group character correspond to ϕ, see [7, Theorem 23.7]. It is easy to see that $\rho \in L^\infty(G)$. Since G is compact, $L^\infty(G) \subseteq L^1(G)$. Then $\rho \in L^1(G)$. Also, since $\rho * f = f * \rho = \phi(f)\rho$ for every $f \in L^1(G)$. One can easily see that ρ is an idempotent in $L^1(G)$. Now by similar argument as in [13, Proposition 2.6], one can easily see that A is approximately biprojective.

Conversely, it is well-known that $L^1(G)$ is biprojective if and only if G is compact. Now apply [13, Proposition 2.4], to complete the proof. □

We recall that a Banach algebra A is left character contractible, if A is left ϕ-contractible for every $\phi \in \Delta(A) \cup \{0\}$, for more information on this notion, see [12].

Proposition 2.5. *Let G be a locally compact group. Then the followings are equivalent:*

(i) *$L^1(G) \otimes_p M(G)$ is biprojective;*
(ii) *$L^1(G) \otimes_p M(G)$ is approximately biprojective;*
(iii) *G is finite.*

Proof. (i)⇒(ii) is clear.

(ii)⇒(iii) Suppose that $L^1(G) \otimes_p M(G)$ is approximately biprojective. Since $M(G)$ is unital, Proposition 2.4 shows that $M(G)$ is approximately biprojective. So by Theorem 1.1 $M(G)$ is left ϕ-contractible for every $\phi \in \Delta(M(G))$. Also by [12, Proposition 3.4] $M(G)$ is left 0-contractible. Hence $M(G)$ is left character contractible. Therefore by [12, Corollary 6.2], G is finite.

(iii)⇒(i) is clear. □

Proposition 2.6. *Let G be an amenable locally compact group. If $L^1(G) \otimes_p \mathcal{A}(G)$ is approximately biprojective, then G is finite.*

Proof. It is well-known that $L^1(G)$ has a bounded approximate identity and by Leptin's theorem amenability of G implies that $\mathcal{A}(G)$ has a bounded approximate identity, see [16, Theorem 7.1.3]. Therefore $L^1(G) \otimes_p \mathcal{A}(G)$ has a bounded approximate identity. Suppose that $L^1(G) \otimes_p \mathcal{A}(G)$ is approximately biprojective. Then by Theorem 1.1, $L^1(G) \otimes_p \mathcal{A}(G)$ is left $\phi \otimes \psi$-contractible for every $\phi \in \Delta(L^1(G))$ and $\psi \in \Delta(\mathcal{A}(G))$. Now by [12, Theorem 3.14], $L^1(G)$ is left ϕ-contractible and $\mathcal{A}(G)$ is left ψ-contractible. By [12, Proposition 6.6], G is discrete and by [12, Proposition 6.1] G is compact, therefore G must be finite. □

3. ϕ-BIFLATNESS

In [17], the authors studied ϕ-biflatness of group algebras. In this section we continue the study of ϕ-biflatness of Segal algebras and the second duals of group algebras.

Theorem 3.1. *Let A be a Banach algebra with a left approximate identity and let $\phi \in \Delta(A)$. If A is ϕ-biflat, then A is left ϕ-amenable.*

Proof. Let A be a ϕ-biflat Banach algebra. Then there exists a bounded A-bimodule morphism $\rho : A \to (A \otimes_p A)^{**}$ such that $\tilde{\phi} \circ \pi_A^{**} \circ \rho(a) = \phi(a)$ for every $a \in A$. Set $g = (id_A \otimes \overline{\phi})^{**} \circ (id_A \otimes q)^{**} \circ \rho : A \to (A \otimes_p \mathbb{C})^{**}$, where $L = \ker \phi$, $q : A \to \frac{A}{L}$ is the quotient map and $\overline{\phi} : \frac{A}{L} \to \mathbb{C}$ is a character defined by

$\overline{\phi}(a+L)=\phi(a)$ for every $a\in A$. We see that g is a bounded left A-module morphism. We claim that $g(l)=0$ for every $l\in L$. Since A has a left approximate identity, $\overline{AL}=L$. Then for each $l\in L$ there exist sequences $(a_n)\subseteq A$ and $(l_n)\subseteq L$ such that $a_nl_n\to l$. For $b\in L$, define a map $R_b:A\to L$ by $R_b(a)=ab$ for every $a\in A$. Since $q\circ R_{l_n}=0$, we have

$$\begin{aligned}
g(l)=(id_A\otimes\overline{\phi})^{**}\circ(id_A\otimes q)^{**}(\rho(l)) &= \lim_n(id_A\otimes\overline{\phi})^{**}\circ(id_A\otimes q)^{**}(\rho(a_nl_n))\\
&=\lim_n(id_A\otimes\overline{\phi})^{**}\circ(id_A\otimes q)^{**}(\rho(a_n)\cdot l_n)\\
&=\lim_n(id_A\otimes\overline{\phi})^{**}\circ(id_A\otimes q)^{**}\circ(id_A\otimes R_{l_n})^{**}(\rho(a_n))\\
&=\lim_n((id_A\otimes\overline{\phi})\circ(id_A\otimes q)\circ(id_A\otimes R_{l_n}))^{**}(\rho(a_n))\\
&=\lim_n((id_A\otimes\overline{\phi})\circ(id_A\otimes(q\circ R_{l_n}))^{**}(\rho(a_n))=0.
\end{aligned}$$

Therefore g induce a map $\overline{g}:\frac{A}{L}\to(A\otimes_p\mathbb{C})^{**}$ which is defined by $\overline{g}(a+L)=g(a)$ for all $a\in A$. It is easy to see that $\overline{g}$ is a bounded left A-module morphism. Pick a_0 in A such that $\phi(a_0)=1$. We denote $\lambda:A\otimes_p\mathbb{C}\to A$ for a map which is specified by $\lambda(a\otimes z)=az$ for every $a\in A$ and $z\in\mathbb{C}$. Set $m=\lambda^{**}\circ\overline{g}(a_0+L)\in A^{**}$, we claim that $am=\phi(a)m$ and $\tilde{\phi}(m)=1$ for every $a\in A$. Since λ^{**} is a left A-module morphism and also since $aa_0+L=\phi(a)a_0+L$, we have

$$am=a\lambda^{**}\circ\overline{g}(a_0+L)=\lambda^{**}\circ\overline{g}(aa_0+L)=\lambda^{**}\circ\overline{g}(\phi(a)a_0+L)=\phi(a)\lambda^{**}\circ\overline{g}(a_0+L)=\phi(a)m \tag{3.1}$$

for every $a\in A$. Since $\rho(a_0)\in(A\otimes_p A)^{**}$, by Goldestine's theorem there exists a net (a_α) in $A\otimes_p A$ such that $a_\alpha\xrightarrow{w^*}\rho(a_0)$. So

$$\begin{aligned}
\tilde{\phi}(m)=m(\phi)=[\lambda^{**}\circ\overline{g}(a_0+L)](\phi) &= [\lambda^{**}\circ g(a_0)](\phi)\\
&=[\lambda^{**}\circ(id_A\otimes\overline{\phi})^{**}\circ(id_A\otimes q)^{**}(\rho(a_0))](\phi)\\
&=[(\lambda\circ(id_A\otimes\overline{\phi})\circ(id_A\otimes q))^{**}(\rho(a_0))](\phi)\\
&=[w^*-\lim(\lambda\circ(id_A\otimes\overline{\phi})\circ(id_A\otimes q))^{**}(a_\alpha))](\phi)\\
&=\lim(\lambda\circ(id_A\otimes\overline{\phi})\circ(id_A\otimes q))^{**}(a_\alpha)(\phi)\\
&=\lim(\lambda\circ(id_A\otimes\overline{\phi})\circ(id_A\otimes q)(a_\alpha)(\phi)\\
&=\lim\phi\circ\lambda\circ(id_A\otimes\overline{\phi})\circ(id_A\otimes q)(a_\alpha)\\
&=\lim\phi\circ\pi_A(a_\alpha).
\end{aligned} \tag{3.2}$$

On the other hand since $a_\alpha\xrightarrow{w^*}\rho(a_0)$, the w^*-continuity of π_A^{**} implies that

$$\pi_A(a_\alpha)=\pi_A^{**}(a_\alpha)\xrightarrow{w^*}\pi_A^{**}(\rho(a_0)).$$

Thus

$$\phi(\pi_A(a_\alpha))=\pi_A(a_\alpha)(\phi)=\pi_A^{**}(a_\alpha)(\phi)\to\pi_A^{**}(\rho(a_0))(\phi)=\tilde{\phi}\circ\pi_A^{**}(\rho(a_0))=1. \tag{3.3}$$

We see that from (3.2) and (3.3), $\tilde{\phi}(m)=1$. Combine this result with (3.1), implies that A is left ϕ-amenable. □

Corollary 3.2. *If $S(G)$ is ϕ-biflat. Then G is amenable*

Proof. Since every Segal algebra has a left approximate identity, by the previous Theorem $S(G)$ is left ϕ-amenable. Then [1, Corollary 3.4] implies that G is amenable. □

We show that the converse of Theorem 3.1 is also valid for symmetric Segal algebras.

Proposition 3.3. *Let G be a locally compact group, and $S(G)$ be a symmetric Segal algebra on G. Then the followings are equivallent*

(i) *G is amenable,*
(ii) *$S(G)$ is ϕ-biflat,*
(iii) *$S(G)$ is left ϕ-amenable.*

Proof. (i)⇒(ii) Let G be an amenable group. Then $L^1(G)$ is amenable. So there exists a bounded net (m_α) in $L^1(G)\otimes_p L^1(G)$ such that $a\cdot m_\alpha - m_\alpha\cdot a\to 0$ and $\pi_{L^1(G)}(m_\alpha)a\to a$ for every $a\in L^1(G)$. It is easy to see that $\phi\circ\pi_{L^1(G)}(m_\alpha)\to 1$ for every $\phi\in\Delta(L^1(G))$. Fix $\phi\in\Delta(L^1(G))$. Define a map $R: L^1(G)\otimes_p L^1(G)\to L^1(G)$ by $R(a\otimes b)=\phi(b)a$ and set $L: L^1(G)\otimes_p L^1(G)\to L^1(G)$ for a map which is specified by $L(a\otimes b)=\phi(a)b$ for every $a,b\in L^1(G)$. It is easy to see that L and R are bounded linear maps which satisfy

$$L(m\cdot a)=L(m)*a,\quad L(a\cdot m)=\phi(a)L(m)\quad (a\in L^1(G), m\in L^1(G)\otimes_p L^1(G))$$

and

$$R(a\cdot m)=a*R(m)\quad R(m\cdot a)=\phi(a)R(m)\quad (a\in L^1(G), m\in L^1(G)\otimes_p L^1(G)).$$

Thus

$$L(m_\alpha)*a-\phi(a)L(m_\alpha)=L(m_\alpha\cdot a-a\cdot m_\alpha)\to 0,$$

similarly we have $a*R(m_\alpha)-\phi(a)R(m_\alpha)\to 0$ for every $a\in L^1(G)$. Since

$$\phi\circ L=\phi\circ R=\phi\circ\pi_{L^1(G)},$$

it is easy to see that

$$\phi\circ L(m_\alpha)=\phi\circ R(m_\alpha)=\phi\circ\pi_{L^1(G)}(m_\alpha)\to 1.$$

Pick an element i_0 in $S(G)$ such that $\phi(i_0)=1$. Set $n_\alpha=R(m_\alpha)i_0\otimes i_0L(m_\alpha)$ for every α. Since $(L(m_\alpha))$ and $(R(m_\alpha))$ are bounded nets in $L^1(G)$ and since $S(G)$ is an ideal of $L^1(G)$, we see that (n_α) is a bounded net in $S(G)\otimes_p S(G)$. Also

$$\begin{aligned}||a\cdot n_\alpha-n_\alpha\cdot a||_{S\otimes_p S}&=||a\cdot n_\alpha-\phi(a)n_\alpha+\phi(a)n_\alpha-n_\alpha\cdot a||_{S\otimes_p S}\\&=||a\cdot n_\alpha-\phi(a)n_\alpha||_{S\otimes_p S}+||\phi(a)n_\alpha-n_\alpha\cdot a||_{S\otimes_p S}\to 0\quad (a\in S(G))\end{aligned}\tag{3.4}$$

and

$$\phi\circ\pi_{S(G)}(n_\alpha)=\phi(R(m_\alpha)*i_0^2*L(m_\alpha))=\phi(R(m_\alpha))\phi(L(m_\alpha))\to 1.\tag{3.5}$$

Let N be a w^*-cluster point of (n_α) in $(S(G)\otimes_p S(G))^{**}$. Combining (3.4) and (3.5) with the facts

$$a\cdot n_\alpha\xrightarrow{w^*}a\cdot N,\quad n_\alpha\cdot a\xrightarrow{w^*}N\cdot a,\quad \pi^{**}_{S(G)}(n_\alpha)\xrightarrow{w^*}\pi^{**}_{S(G)}(N)\quad (a\in(S(G))$$

we have

$$a\cdot N=N\cdot a,\quad \tilde{\phi}\circ\pi^{**}_{S(G)}(N)=1\quad (a\in(S(G))).$$

Define a map $\rho : S(G) \to (S(G) \otimes_p S(G))^{**}$ by $\rho(a) = a \cdot N$ for every $a \in S(G)$. It is easy to see that ρ is a bounded $S(G)$-bimodule morphism and $\tilde{\phi} \circ \pi^{**}_{S(G)} \circ \rho(a) = \tilde{\phi} \circ \pi^{**}_{S(G)}(a \cdot N) = \phi(a)$, so $S(G)$ is ϕ-biflat.
(ii)⇒(i) is clear by Corollary 3.2.
(iii)⇔(i) is clear by [1, Corollary 3.4]. □

Let A be a Banach algebra and $\phi \in \Delta(A)$. A is called ϕ-inner amenable if there exists an element $m \in A^{**}$ such that $m(f \cdot a) = m(a \cdot f)$ and $\tilde{\phi}(m) = 1$ for every $a \in A$ and $f \in A^*$, see [8]. Note that by [8, Corollay 2.2] every Banach algebra with a bounded approximate identity is ϕ-inner amenable.

Theorem 3.4. *Let A be a ϕ-inner amenable Banach algebra, where $\phi \in \Delta(A)$. If A^{**} is $\tilde{\phi}$-biflat, then A is left ϕ-amenable.*

Proof. Let A^{**} be $\tilde{\phi}$-biflat. Then there exists a bounded A^{**}-bimodule morphism $\rho : A^{**} \to (A^{**} \otimes_p A^{**})^{**}$ such that for every $a \in A^{**}$

$$\tilde{\tilde{\phi}} \circ \pi^{**}_{A^{**}} \circ \rho(a) = \tilde{\phi}(a),$$

where $\tilde{\tilde{\phi}}$ is an extension of $\tilde{\phi}$ on A^{****} as we mentioned in the introduction. Suppose that A is ϕ-inner amenable. Then there exists an element $m \in A^{**}$ such that $m(f \cdot a) = m(a \cdot f)$ and $\tilde{\phi}(m) = 1$ for every $a \in A$ and $f \in A^*$. Set $M = \rho(m)$, since ρ is a bounded A^{**}-bimodule morphism, we have $a \cdot M = M \cdot a$ and $\tilde{\tilde{\phi}} \circ \pi^{**}_{A^{**}}(M) = \tilde{\phi}(m) = 1$ for every $a \in A$.

Now take $\epsilon > 0$ and a finite set $F = \{a_1, ..., a_r\} \subseteq A$, and set

$$\begin{aligned} V =& \{(a_1 \cdot n - n \cdot a_1, ..., a_r \cdot n - n \cdot a_r, \tilde{\phi} \circ \pi_{A^{**}}(n) - 1) : n \in A^{**} \otimes_p A^{**}, ||n|| \leq ||M||\} \\ &\subseteq \prod_{i=1}^{r} (A^{**} \otimes_p A^{**}) \oplus_1 \mathbb{C}. \end{aligned}$$

Then V is a convex set and so the weak and the norm closures of V coincide. But by Goldestine's theorem there exists a net $(n_\alpha) \subseteq A^{**} \otimes_p A^{**}$ such that $n_\alpha \xrightarrow{w^*} M$ and $||n_\alpha|| \leq ||M||$. So for every $a \in F$ we have $a \cdot n_\alpha - n_\alpha \cdot a \xrightarrow{w} 0$ and $|\tilde{\phi} \circ \pi_{A^{**}}(n_\alpha) - 1| \to 0$ which shows that $(0, 0, ..., 0)$ is a $||\cdot||$-cluster point of V. Thus there exists an element $n_{(F,\epsilon)}$ in $A^{**} \otimes_p A^{**}$ such that

$$||a_i \cdot n_{(F,\epsilon)} - n_{(F,\epsilon)} \cdot a_i|| < \epsilon, \quad |\tilde{\phi} \circ \pi_{A^{**}}(n_{(F,\epsilon)}) - 1| < \epsilon \tag{3.6}$$

for every $i \in \{1, 2, \ldots, r\}$. Now we consider a directed set

$$\Delta = \{(F, \epsilon) : F \text{ is a finite subset of } A, \epsilon > 0\},$$

with the following order

$$(F, \epsilon) \leq (F', \epsilon') \Longrightarrow F \subseteq F', \quad \epsilon \geq \epsilon'.$$

So the equation (3.6) follows that there exists a bounded net $(n_{(F,\epsilon)})_{(F,\epsilon)\in\Delta}$ in $A^{**} \otimes_p A^{**}$ such that

$$a \cdot n_{(F,\epsilon)} - n_{(F,\epsilon)} \cdot a \to 0, \quad \tilde{\phi} \circ \pi_{A^{**}}(n_{(F,\epsilon)}) \to 1$$

for every $a \in A$. By [4, Lemma 1.7] there exists a bounded linear map $\psi : A^{**} \otimes_p A^{**} \to (A \otimes_p A)^{**}$ such that for $a, b \in A$ and $m \in A^{**} \otimes_p A^{**}$, the following holds

(i) $\psi(a \otimes b) = a \otimes b$,
(ii) $\psi(m) \cdot a = \psi(m \cdot a), \qquad a \cdot \psi(m) = \psi(a \cdot m)$,
(iii) $\pi^{**}_A(\psi(m)) = \pi_{A^{**}}(m)$.

Define $\xi_{(F,\epsilon)} = \psi(n_{(F,\epsilon)})$ which is a net in $(A \otimes_p A)^{**}$ and by the previous properties of ψ it satisfies

$$a \cdot \xi_{(F,\epsilon)} - \xi_{(F,\epsilon)} \cdot a \to 0, \quad \tilde{\phi} \circ \pi_A^{**}(\xi_{(F,\epsilon)}) \to 1 \quad (a \in A).$$

Now by applying a similar method as we obtained a net from M at the beginning of the proof, one can obtain a bounded net $(\gamma_{(F,\epsilon)})_{(F,\epsilon)\in\Delta}$ related to $\xi_{(F,\epsilon)}$ in $A \otimes_p A$ such that

$$a \cdot \gamma_{(F,\epsilon)} - \gamma_{(F,\epsilon)} \cdot a \to 0, \quad \phi \circ \pi_A(\gamma_{(F,\epsilon)}) \to 1 \quad (a \in A).$$

Now define $T : A \otimes_p A \to A$ by $T(a \otimes b) = \phi(b)a$ for every a and b in A. It is easy to see that T is a bounded linear map with the following properties

$$T(a \cdot m) = aT(m), \quad T(m \cdot a) = \phi(a)T(m) \qquad (m \in A \otimes_p A, \quad a \in A).$$

Define $\nu_{(F,\epsilon)} = T(\gamma_{(F,\epsilon)})$, it is easy to see that $\nu_{(F,\epsilon)}$ is a bounded net and

$$a\nu_{(F,\epsilon)} - \phi(a)\nu_{(F,\epsilon)} \to 0, \quad \phi \circ T(\nu_{(F,\epsilon)}) = \phi \circ \pi_A(\gamma_{(F,\epsilon)}) \to 1 \quad (a \in A).$$

Therefore by [9, Theorem 1.4] A is left ϕ-amenable. □

Corollary 3.5. *Let G be a locally compact group. If $L^1(G)^{**}$ is $\tilde{\phi}$-biflat, then G is amenable.*

Proof. Since $L^1(G)$ has a bounded approximate identity, $L^1(G)$ is ϕ-inner amenable. Thus by Theorem 3.4, $L^1(G)$ is left ϕ-amenable. Now by [1, Corollary 3.4] G is amenable. □

Corollary 3.6. *Let G be a locally compact group and $\phi, \psi \in \Delta(L^1(G))$. If $(M^1(G) \otimes_p L^1(G))^{**}$ is $\widetilde{\phi \otimes \psi}$-biflat, then G is amenable.*

Proof. We note that $M(G) \otimes_p L^1(G)$ has a bounded approximate identity and so it is ϕ-inner amenable. Now by Theorem 3.4, $M(G) \otimes_p L^1(G)$ is left $\phi \otimes \psi$-amenable, where $\phi, \psi \in \Delta(L^1(G))$. Hence by [9, Theorem 3.3], $L^1(G)$ is left ϕ-amenable, hence G is amenable. □

References

[1] M. Alaghmandan, R. Nasr Isfahani and M. Nemati; *Character amenability and contractibility of abstract Segal algebras*, Bull. Aust. Math. Soc, **82** (2010) 274-281.

[2] Y. Choi, F. Ghahramani, Y. Zhang *Approximate and pseudo-amenability of various classes of Banach algebras*, J. Funct. Anal. **256** (2009) 3158-3191.

[3] P. Eymard; *Làlgèbre de Fourier d́un groupe localement compact*, Bull. Soc. Math. France **92** (1964), 181-236.

[4] F. Ghahramani, R. J. Loy and G. A. Willis; *Amenability and weak amenability of second conjugate Banach algebras*, Proc. Amer. Math. Soc. **124** (1996), 1489-1497

[5] F. Ghahramani, Y. Zhang; *Pseudo-amenable and pseudo-contractible Banach algebras*, Math. Proc. Cambridge Philos. Soc. **142** (2007) 111-123.

[6] A. Ya. Helemskii; *The homology of Banach and topological algebras*, Kluwer, Academic Press, Dordrecht, 1989.

[7] E. Hewitt and K. Ross; *Abstract Harmonic Analysis I*, Die Grundlehren der Mathematischen Wissenschaften, 115, Springer-Verlag, Berlin, (1963).

[8] A. Jabbari, T. Mehdi Abad and M. Zaman Abadi; *On ϕ-inner amenable Banach algebras*, Colloq. Math. vol **122** (2011) 1-10.

[9] E. Kaniuth, A. T. Lau and J. Pym; *On ϕ-amenablity of Banach algebras*, Math. Proc. Cambridge Philos. Soc. **144** (2008) 85-96.

[10] J. L. Kelley; *General topology*, D. Van Nostrand Company, Inc., Toronto-New York- London, 1955.

[11] E. Kotzmann and H. Rindler; *Segal algebras on non-abelian groups*, Trans. Amer. Math. Soc. **237** (1978), 271-281.

[12] R. Nasr Isfahani and S. Soltani Renani; *Character contractibility of Banach algebras and homological properties of Banach modules*, Studia Math. **202** (2011) 205-225.

[13] P. Ramsden; *Biflatness of semigroup algebras*, Semigroup Forum **79**, (2009) 515-530.

[14] H. Reiter; L^1*-algebras and Segal Algebras*, Lecture Notes in Mathematics **231** (Springer, 1971).

[15] H. Reiter, J.D. Stegeman, Classical Harmonic Analysis and Locally Compact Groups, 2nd ed., London Math. Soc. Monogr., vol. 22, Oxford Univ. Press, New York, 2000.

[16] V. Runde; *Lectures on Amenability*, (Springer, New York, 2002).

[17] A. Sahami and A. Pourabbas; *On ϕ-biflat and ϕ-biprojective Banach algebras*, Bull. Belg. Math. Soc. Simon Stevin, **20**(2013) 789-801.

[18] A. Sahami and A. Pourabbas; *Approximate biprojectivity of certain semigroup algebras*, see arXiv:1409.7515v1 [math.FA] 26 Sep 2014.

[19] E. Samei, N. Spronk and R. Stokke; *Biflatness and pseudo-amenability of Segal algebras*, Canad. J. Math. **62** (2010), 845-869.

[20] Y. Zhang; *Nilpotent ideals in a class of Banach algebras*, Proc. Amer. Math. Soc. **127** (11) (1999), 3237-3242.

Faculty of Mathematics and Computer Science, Amirkabir University of Technology, 424 Hafez Avenue, 15914 Tehran, Iran.

E-mail address: amir.sahami@aut.ac.ir

E-mail address: arpabbas@aut.ac.ir